



Università degli Studi di Siena 1240-2010: 770 years of history





Siena, Policlinico Universitario "Santa Maria alle Scotte"




Endometriosis: symptoms



Endometriosis and infertility: possible mechanisms

- Follicular environment and ovulation
- Changes in peritoneal fluid
- ❖ Adhesions between peritoneum and pelvic organs
- Eutopic endometrium abnormalities
- Implantation disorders





Progesterone






Growth factors

Cytokines Chemokines


Free radicals (Oxidative stress)

Prostaglandins




Inflammation

Endometriosis: tissues studied




Eutopic endometrium



But also:

- Gastrointestinal lesions
- Urinary tract

Eutopic and ectopic endometrium: hormone receptors, factors and enzymes




Neuroendocrine factors in endometriosis




In women with endometriosis, there is abnormal sprouting of nociceptors in the endometrium and in peritoneal endometriotic lesions


Nerve fibres



Jabbour et al. SGI 2010



Neuroendocrine factors in endometriosis



Substance P

VIP

Synaptophysin (SYN)


Neuropeptide Y

Calcitonine gene related protein

neuron-specific enolase (NSE)

Bokor et al, 2009 and Wang et al., 2010

Neuroendocrine system is localized in various organs and tissues



Endometrium and neuroendocrine factors

Prolactin


CRH


Urocortins

Relaxin

Substance P


Vasoactive Intestinal peptide








CRH is a neuropeptide originally localized in central nervous system




CRH-family

	amino acids		% CRH sequence homology
CRH	41		
urocortin	40		42
Vaughan et al., Nature 1995			
urocortin 2 (orstresscopin related peption	de) 38		31
Reyes et al., PNAS U S A. 20	001		
urocortin 3 (pr stresscopin)	38		26
Lewis et al., PNAS U S A. 20 <mark>01</mark>			

CRH- Receptors affinity



Human endometrium express and localize



CRH

Urocortin


Urocortin 2

Urocortin 3

CRH-R1

CRH-R2


Endometrial CRH/Ucn and menstrual cycle



CRH mRNA expression is higher during secretory phase (Zoumakis et al. 2001)

Ucn mRNA expression is higher during secretory phase (Torricelli et al, Hum Reprod 2007)

Endometrial cells and CRH receptors



Endometrial cells express
CRH-R1 mRNA in both phases
of menstrual cycle

Proliferative phase


Secretory phase


Endometrial cells express CRH-R2 mRNA in both phases of menstrual cycle

Karteris et al., JCE&M 2004

Di Blasio et al., JCE&M 1997

CRH stimulates decidualization in endometrial stromal cells







Zoumakis et al., Mol Hum Rep 2000

Ferrari et al., J Steroid Biochem Mol Biol.1995

Ucn stimulates decidualization in endometrial stromal cells



Ucn and prolactin concentration Increase during in vitro decidualization



Ucn stimulates decidualization process


CRH/CRH-R1 are involved in embryo implantation

Mouse model


Antalarmin


Block of CRH-R1

Makrigiannakis et al., Am J Reprod Immun 2004




Implantation failure 70%







Kalantaridou et al., Crit Rev in Clinic Lab Sci 2007

Endometriosis and CRH




Endometriosis and Ucn







Ectopic tissue express Ucn mRNA

Urocortin levels in ovarian endometriosis




Endometrium and Ucn







Ucn level detects 88% of cases of endometrioma with a specificity of 90%


Eutopic endometrium in endometriosis: impaired CRH and Ucn mRNA expression






Eutopic endometrium in endometriosis: role of CRH and Ucn in *in vitro* decidualization








Submitted 2010

Role of CRH and Ucn in endometriosis


CRH and Ucn showed aberrant cycle-related changes in endometriosis

The lack of increase of CRH and Ucn during secretory phase may be involved in a deranged decidualization


Endometriotic cells show a deranged expression of CRH-R1




CRH-family





Endometrium and Ucn2/Ucn3



Endometrium and Ucn2/Ucn3









Endometriosis and Ucn2/Ucn3

Ectopic lesions express and localize Ucn2 and Ucn3 mRNA

UCN2



UCN3




UCN2 / HPRT

Relative mRNA expression


80.00

10.00


Ectopic



Ucn2 /Ucn3 in endometrium of endometriotic women



Ucn2 mRNA expression showed the highest expression in the late secretory phase



Ucn3 mRNA expression did not change during endometrial cycle

Urocortin2 and inflammation



Urocortin 2



Proinflammatory action

Moss et al. Gut 2007

Antiinflammatory action

Proinflammatory action

Submitted 2010

Urocortin 3 and inflammation



Urocortin 3




Submitted 2010

Saruta et al., Peptides 2005

Antiinflammatory
Action



Effect of Ucn2 in cultured endometrial stromal cells on cytokines secretion




Ucn2 treatment increased TNF-α release in a dose dependent manner

Astressin 2b reversed Ucn2 effects

Submitted 2010



Effect of Ucn3 in cultured endometrial stromal cells on cytokines secretion




Ucn3 treatment increased IL-4 release


Such effect was reversed by astressin 2b

Submitted 2010

Neuroendocrine elements in endometriosis



Neuroendocrine elements in endometriosis



Acknowledgements



Clinicians

Laboratory group



Key points of established knowledge

- 1. Endometrial dysfunction is a cause of infertility in endometriosis
- 2. Increased abortion rate in endometriosis is related to endometrial dysfunction
- 3. Estrogen/Progesterone act on endometrial functions through locally produced growth factors/peptides and the alteration of this paracrine actions occurs in endometriosis
- 4. Endometrial cells implant in peritoneum is driven by immunomodulatory factors, facilitating neoangiogenesis and decreasing apoptotic process
- 5. Ectopic endometrium expresses some of the growth factors/peptides functionally produced by eutopic endometrium, with an overfunction of proinflammatory pathways

Unresolved areas

- A. Increased estrogen activity or progesterone resistance remain to be demonstrated in a large scale study by gene mutations / polimorphism evaluations
- B. Has ectopic endometrium identical characteristics of the eutopic tissue?
- C. The use of endometrial cells or factors for the noninvasive diagnosis of endometriosis is under investigation
- D. The endometrial action of estroprogestins in the treatment of endometriosis is not yet defined

Further research questions (1)

Epidemiology: it is not clear how is really endometriosis diffuse in the

Developing Countries

Classification: is deep infiltrating endometriosis a separate disease or just

a developmental stage?

<u>Pathogenesis</u>: How critical are the serum estrogen/progesterone level

changes in creating the endometrial substrate for

endometriosis?

How relevant are the epigenetic influences on

estrogen/progesterone receptors?

Is infertility related to specific endometrial pathways

dysfunction?

<u>Diagnosis</u>: Endometrial cytology may have a future for the diagnosis

of endometriosis? Urinary or serum markers have

sufficient specificity? Does uterine doppler analysis reveal

some specific changes?

Further research questions (2)

Treatment: Do surgical procedures contribute to spread

endometriotic cells?

Does it exist an optionally timing for the surgical

treatment of endometriosis?

Is induction of ovulation a treatment at risk for women

with endometriosis?

How to prevent the recurrency?

Women's Health: How is the gestational outcome?

How is he climateric period?

How is the postmenopausal period in Women with

endometriosis?